Check List the journal of biodiversity data
NOTES ON GEOGRAPHIC DISTRIBUTION

Occurrence of chitons (Mollusca, Polyplacophora) in estuaries: first records in Brazil

Davi J. Araújo de Lima[1], Rafaela Camargo Maia[2], Jaime Alberto Jardim[3]

1 *Programa de Pós-graduação em Biologia Animal. Universidade Federal dos Vales do Jequitinhonha e Mucuri, Alto da Jacuba, Diamantina, MG, Brazil*
2 *Laboratório Ecologia de Manguezais, Instituto Federal do Ceará, Avenida Desembargador Armando de Sales Louzada, s/n, Monsenhor Edson Magalhães, Acaraú, CE, Brazil*
3 *Museu de Zoologia da Universidade de São Paulo, Avenida Nazareth, 481, Ipiranga, São Paulo, SP, Brazil*

Corresponding author: *Davi J. Araújo de Lima (davijoseh89@gmail.com)*

Abstract. Chitons are molluscs with scarce world records in estuaries. In the present study, we provide the first records of two species of Polyplacophora in the Coreaú River estuary, Camocim, Ceará, Brazil: *Ischnochiton striolatus* and *Ischnoplax pectinata*. The data shows these animals thrive in estuary coastal regions. The taxonomic data together with the ecological data presented in this study, it is possible to conclude that this research represents a significant milestone in understanding the biodiversity of Brazilian estuarine areas.

Key words. Chitonida, mangrove, northeastern Brazil, semi-arid coast

Lima DJA, Maia RC, Jardim JA (2024) Occurrence of chitons (Mollusca, Polyplacophora) in estuaries: first records in Brazil. Check List 20 (4): 908–914. https://doi.org/10.15560/20.4.908

INTRODUCTION

Estuaries are coastal bodies of water that have a free connection to the open sea and dilution with freshwater draining from off the land. Variation in salinity and passage of nutrients and sediments brought by water from inland to coastal regions is common (Silva 2000; Dias et al. 2016), forming diverse habitats for benthic organisms throughout their life cycle (Menegotto et al. 2019). Knowledge of the distribution of marine species in an estuary can be useful for assessing changes in benthic habitats resulting from sea-level rise attributed to global warming (Shaw et al. 2009).

The class Polyplacophora is restricted to marine environments, where they live at depths down to 8,000 m (Kaas and Jones 1998) and shallower, coastal environments, such as rocky beaches, tide pools, and reefs (Eernisse and Clark 1994; Guimarães et al. 2008; Schwabe 2010; Veiga et al. 2016). Few studies have reported the occurrence of chitons in estuaries (Figure 1). Chalmer and Hodgkin (1976) first reported chitons in an estuary, in the Swan River estuary of Western Australia; they found *Acanthochitona* sp. (Acanthochitonidae) and a lepidopleurid of uncertain genus in this temperate estuary. In the same region, Shaw et al. (2009) found the following species: *Ischnochiton cariosus* Carpenter, 1879 (Ischnochitonidae), *Ischnochiton contractus* (Reeve, 1847) (Ischnochitonidae), *Cryptoplax striata* (Lamarck, 1819) (Cryptoplacidae), and *Acanthochitona bednalli* (Pilsbry, 1894).

Giberto et al. (2004) recorded the occurrence of *Chaetopleura isabellei* (d'Orbigny, 1841) (Chaetopleuridae) in the estuary of the Rio de la Plata, between Argentina and Uruguay. Wolff (2005) collected *Leptochiton cancellatus* (G.B. Sowerby II, 1840) (Leptochitonidae) in the Oosterschelde estuary, Holland. Arias and Anadón (2013) indicated the presence of *Tonicia atrata* (G.B. Sowerby II, 1839) (Chitonidae) in the Eo and Avilés estuaries (Spain), and in the Sado estuary (Portugal). There are no records of chitons in estuaries from other regions; however, polyplacophorans are little studied when compared to other classes of molluscs (Correia et al. 2015). In the present study, we document the first occurrence of chitons in an estuary in Brazil.

OPEN ACCESS

Academic editor: Juan Francisco Araya
Received: 26 January 2024
Accepted: 21 May 2024
Published: 31 July 2024

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

METHODS

The sampling of chitons was carried out at low spring tide on the rocky shore of the lower estuary of the Coreaú River, which is in the municipality of Camocim, Ceará, northeastern Brazil. The estuary is on the



Figure 1. New record (light grey circle) and previous records (dark grey circles) of Polyplacophora in estuaries.

equatorial coast of the semi-arid region of Brazil (Morais et al. 2020), where the average annual rainfall is 1,114 mm (Climate-data.org n.d.).

The Coreaú River originates at the confluence of Jatobá and Caiçara streams, at the foot of the Serra da Ibiapaba. It flows for 167.5 km before entering the Atlantic Ocean. The estuary of this river features sediment rich in organic matter sourced from nearby mangroves along the shores, along with the presence of silico-ferruginous rocks and detritus and sandstone reefs (Morais et al. 2006).

Bimonthly, between March 2017 and March 2019, three sampling points were selected along the estuarine strip of the Coreaú River. The points were spaced 2 km apart. At each point, the distance to the low tide line was 30 m^2 in all intertidal zones, and for each point three random quadrants with an area of 5 m^2 each were delineated. In each quadrant, chitons were collected through the active search method, for 10 min with the sampling effort of four observers. The ambient air temperature was measured at each sampling point with a thermo-hygrometer, and salinity was measured with a refractometer. Rainfall data were extracted from the Fundação Cearense de Meteorologia e Recursos Hídricos, Funceme (2017–2019).

In the laboratory, animals were fixed and preserved in 70% alcohol. The taxonomic information for diagnosis and identification of the material was based on the literature (Righi 1970, 1971; Kaas and Belle 1985, 1990, 1994; Simone and Jardim 2009). Measurements of length, spanning from the cephalic to the anal valve, were acquired from diagnostic individuals utilizing digital calipers with 0.01 mm precision.

Individuals were classified into three size categories, as defined by (Iredale et al. 1924; Schwabe 2010): small individuals, where the length was ≤15 mm; medium individuals, where the length ranged from >15 mm to ≤30 mm; and large individuals, with a length >30 mm. To analyze sampling effort efficiency, an accumulation curve based on the monthly sampling effort (Jackknife) was performed. After analysis in the laboratory, a few specimens of each species were selected for vouchers and deposited in the Collection Malacological Prof. Henry Ramos Matthews (**CMPHRM**) from the Institute of Marine Sciences at the Federal University of Ceará.

Only specimens deposited in the CMPHRM collection were used in describing diagnostic characters. The other, non-vouchered specimens were only considered in our study of abiotic factors and abundance. We observed and compared variations in abiotic factors, such as temperature and salinity, and abundance during the sampling period.

RESULTS

In our study, we found 981 individuals, which belong two species: *Ischnochiton striolatus* (Gray, 1828) with 962 individuals, and *Ischnoplax pectinata* (G.B. Sowerby II, 1840) with 19 individuals. Figure 2 shows living animals of both these species on their respective substrates.

Family Ischnochitonidae

***Ischnochiton striolatus* (Gray, 1828)**
Figure 2A

Material examined. BRAZIL –Ceará • Camocim, Coreaú River estuary; 02°52.125′S, 040°51.425′W; on a thick, small rock in a rocky area of a sandstone reef within tide pools; D.J. Lima & R.C. Maia leg.; 08.VII.2017; 6 spec., MPHRM 5144.

Diagnostic characters. Animal elongate-oval; small (length 9.20–3.52 mm); tegmentum slightly grayish–yellowish white, with mixed colors, spotted with beige. Dorsal valve surface striate.

Family Callistoplacidae

***Ischnoplax pectinata* (G.B. Sowerby II, 1840)**
Figure 2B

Material examined. BRAZIL – Ceará • Camocim, Coreaú River estuary; 02°53.143′S, 040°50.742′W; on a thick, small rock in a rocky area of a sandstone reef within tide pools; D.J. Lima & R.C. Maia leg.; 08.VII.2017; 2 spec., CMPHRM 5143.

Diagnostic characters. Animal very elongate-oval; small to large (length 9.75–17.17 mm); tegmentum colour varying from beige (sometimes greyish) to greenish. Dorsal valve sculpture with radial ribs on the head valve has radial ribs; median valves have ribs on margins in lateral and pleural areas; central region of median valves with a smooth jugal line; terminal valve with ribs. Anteromucronal area (before the apex) concave and with a prominent apex.

Ecological aspects. The accumulation curve based on the bimonthly sampling effort indicated an estimate of the richness of five chiton species (Jackknife = 4.99 ± 0.59), with a collection efficiency coefficient of 80.16%. Throughout the sampling period, the ambient air temperature ranged from 22.5 °C on September 18 to a peak of 41 °C on March 2017 (with an average temperature of 31 °C ± 0.31 °C). Concurrently, salinity levels ranged between 8.20 ppt. on March 19 and 43 ppt. on September 2018 (with an average salinity of



Figure 2. Species of Polyplacophora in the estuary of the Coreaú River, Ceará, Brazil. **A.** *Ichnochiton striolatus*. **B.** *Ischnoplax pectinata*.

32.5 ppt. ± 0.78 ppt.). The greatest abundance of *Ischnochiton striolatus* was observed on July 17, with 158 individuals, *Ischnoplax pectinata* was most numerous on May 2018, with seven individuals (Figure 3). The average rainfall during the study ws 106.54 mm ± 16.97, and greatest rainfall was on March 2019 collection (Figure 4).

DISCUSSION

We record for the first time the occurrence of two species of Polyplachophora in a Brazilian estuary. Until now, chitons had only been observed in estuaries in temperate regions (Giberto et al. 2004; Wolff 2005; Arias and Anadón 2013).

Ischnochiton striolatus and *Ischnoplax pectinata* both occur along the coast of the Western Atlantic Ocean, from North Carolina, USA, to Santa Catarina, Brazil (Simone and Jardim 2009; Jardim 2015; Matthews-Cascon and Menezes 2019). In the state of Ceará, Veras et al. (2013) found *Ischnochiton striolatus* and *Ischnoplax pectinata* living on sandstone reefs at Pacheco Beach. Rocha-Barreira et al. (2017) also reported the occurrence of *Ischnochiton striolatus* on rocky beaches and marine angiosperms banks, and they noted that the occurrence of *Ischnoplax pectinata* was restricted to sandstone reefs. Lopes et al. (2019) documented the occurrence of *I. striolatus* on Picos Beach in Icapuí, where this chiton settled on shells of the gastropod *Turbinella laevigata* Anton, 1838.

From our collections, we identified two species, but the accumulation curve indicates that, with greater temporal and spatial replication, there potentially could be three additional species found. This highlights the effectiveness of our analysis in terms of the methods applied and the sampling duration. However, further studies in similar estuarine areas are needed prove the curve analysis correct.

The variation in salinity at our sampling points revealed that these chitons can tolerate extreme conditions, as evidenced by the lower salinity levels observed on March 19, which was due to a prevailing, seasonal rainy period at that time (Figure 3). The stress caused by reduced salinity in estuaries is the main factor responsible for structuring faunal communities (Nebra et al. 2016). Estuaries in tropical and arid regions have higher salinity than estuaries in temperate regions due to much less freshwater reaching the sea and because of higher insolation (Eyre and Balls 1999). Thus, the Coreaú river estuary is a propitious environment for the occurrence of chitons, as seem by the abundance of specimens in the Coreaú river estuary, even amid seasonal variations.

Similar to the decrease in salinity, rainy seasons are associated with a reduction in the abundance, as evidenced by a decrease in abundance of chitons on March 17–18, May 18, and March 19. Conversely, in other months, except for September 18 and November 18, individuals were more abundant (Figure 4). Our abundance data, along with abiotic factors, do not support more complex ecological analyses; they merely serve to characterize the environment in our study. Variable characteristics present in the Coreaú estuary suggest that habitats there are favorable for the development of a chiton community. This hypothesis agrees with the results obtained by Vidy (2000), who found that estuarine environments represent natural nurseries for marine life.

Other factors, such as soil-texture composition, can also modify the distribution of organisms in an estuary (Boyle 1969; Rostal and Simpson 1988; Grayson and Chapman 2004). Chitons are associated with consolidated substrates, which are uncommon in estuaries (Eernisse and Clark 1994). In the Coreaú River, the presence of sandstone reefs strongly contributes to the establishment of these animals, as these reefs—in areas dominated by soft substrates—provide a solid substrate that allows for the movement of individuals



Figure 3. Sample variation graph based on the abundance results of the species *Ischnochitons striolatus* and *Ischnoplax pectinata*, and abiotic characteristics (salinity and air temperature) in relation to the sampling axis from March 17–19.



Figure 4. Sample variation graph based on the abundance results of the *species Ischnochitons striolatus* and *Ischnoplax pectinata*, and rainfall in relation to the sampling axis from March 17–19.

for feeding and serves as refuges against predators, which results in a greater survival of young and, thus, in a higher density of individuals (Ferreira et al. 2007; Mendonça et al. 2015; Lefcheck et al. 2019).

Our research is a significant in better understanding the biodiversity of Brazilian estuarine areas. The new data represent the first recorded occurrences of chitons in an estuary in the region, contributes to the knowledge of the geographic distribution of these organisms, and elucidates their environmental adaptation to the Rio Coreaú estuary. Additionally, data on seasonal variations in temperature, salinity, and precipitation provides important insights into the environmental factors that may influence the abundance and distribution of these species. Our results are may help support conservation and management measures for estuarine ecosystems and highlight the importance of scientific research for the preservation of Brazilian marine biodiversity.

ACKNOWLEDGEMENTS

We thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (Cnpq) and the Instituto Federal do Ceará for financial and logistical support. The Laboratório de Ecologia de Manguezais and its staff are acknowledged for providing resources. To the Jornal's academic editors and reviewers for the newspaper for improving and textual refinement of this manuscript.

ADDITIONAL INFORMATION

Conflict of interest
The authors declare that no competing interests exist.

Ethical statement
No ethical statement is reported.

Funding
This study was financially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ).

Author contributions
Conceptualization: DJAL, RCM. Funding acquisition: RCM. Investigation: DJAL, RCM, JAJ. Methodology: DJAL, RCM, JAJ. Project administration: RCM. Writing – original draft: DJAL. Writing – review and editing: DJAL, RCM, JAJ.

Author ORCID ids
Davi J. Araújo de Lima https://orcid.org/0000-0002-7959-9508
Rafaela Camargo Maia https://orcid.org/0000-0001-5871-4610
Jaime Alberto Jardim https://orcid.org/000-0002-5511-7172

Data availability
All data that support the findings of this study are available in the main text.

REFERENCES

Arias A, Anadón N (2013) *Tonicia atrata* and *Chiton cumingsii* (Polyplacophora: Chitonidae): first records in European waters. Zootaxa 3626: 593–596. https://doi.org/10.11646/zootaxa.3626.4.14

Boyle PR (1969) The survival of osmotic stress by *Sypharochiton pelliserpentis* (Mollusca: Polyplacophora). The Biological Bulletin 136: 154–166. https://doi.org/10.2307/1539810

Chalmer PN, Hodgkin EP (1976) Benthic faunal changes in a seasonal estuary of south-western australia. Records of the Western Australian Museum 4: 383–410.

Climate-data.org (n.d.) https://pt.climate-data.org/america-do-sul/brasil/ceara/acarau-44118/. Accessed on: 2020-01-27.

Correia MD, Coelho CA, Sovierzoski HH (2015) Polyplacophora (Mollusca) from reef ecosystems and associations with macroalgae on the Coast of Alagoas, northeastern Brazil. Zoologia 32: 289–295. https://doi.org/10.1590/S1984-46702015000400004

Dias FJS, Castro BM, Lacerda LD, Miranda LB, Marins RV (2016) Physical characteristics and discharges of suspended particulate matter at the continent-ocean interface in an estuary located in a semiarid region in northeastern Brazil. Estuarine, Coastal and Shelf Science 180: 258–274. https://doi.org/10.1016/j.ecss.2016.08.006

Eernisse DJ, Clark RN (1994) Polyplacophora. In: Harrison FW, Kohn AJ (Eds.) Microscopic anatomy of invertebrates. New York, United States, 5: 56-110.

Eyre BBP (1999) A comparative study of nutrient behavior along the salinity gradient of tropical and temperate estuaries. Estuaries 22: 313–326. https://doi.org/10.2307/1352987

Ferreira TO, Vidal-Torrado P, Otero XL, Macías F (2007) Are mangrove forest substrates sediments or soils? A case study in southeastern Brazil. Catena 70: 79–91. https://doi.org/10.1016/j.catena.2006.07.006

Funceme (2017–2019) https://chuvas.funceme.br/diario/municipios/maxima/2017/3. Accessed on: 2017-03-17.

Giberto DA, Bremec CS, Acha EM, Mianzan H (2004) Large-scale spatial patterns of benthic assemblages in the SW Atlantic: the Río de la Plata estuary and adjacent shelf waters. Estuarine, Coastal and Shelf Science 61: 1–13. https://doi.org/10.1016/j.ecss.2004.03.015

Grayson JE, Chapman MG (2004) Patterns of distribution and abundance of chitons of the genus Ischnochiton in intertidal boulder fields. Austral Ecology 29: 363–373. https://doi.org/10.1111/j.1442-9993.2004.01375.x

Guimarães IM, Antonio ÍG, Peixoto S, Olivera A (2008) Salinity influence on the survival of mangrove oyster, Crassostrea rhizophorae. Arquivo de Ciência do Mar 41:118–22.

Iredale T, Iredale T, Hull AFB (1924) A monograph of the Australian loricates (Phylum Mollusca – Order Loricata). III. The Australian Zoologist 3: 277–297.

Jardim JA (2015) Revisão taxonômica do gênero *Ischnoplax* (Chitonoidea; Callistoplacidae) do Atlântico Oeste. Master's thesis, Universidade de São Paulo, São Paulo, Brazil, 88 pp. https://doi.org/10.11606/d.38.2016.tde-17062016-205541

Lefcheck JS, Hughes BB, Johnson AJ, Pfirrman BW, Rasher DB, Smyth AR, Williams BL, Beck MW, Orth RJ (2019) Are coastal habitats important nurseries? A meta-analysis. Conservation Letters 12: e12645. https://doi.org/10.1111/conl.12645

Kaas P, Belle RAV (1985) Monograph of Living Chitons: (Mollusca: Polyplacophora). E.J. Brill, Leiden, the Netherlands, 244 pp.

Kaas P, Van Belle RA (1990) Monograph of living chitons: (Mollusca: Polyplacophora). Volume 4. Suborder Ischnochitonina: Ischnochitonidae:Ischnochitoninae (continued), additions to vols. 1, 2 y 3. E.J. Brill, Leiden, the Netherlands, 298 pp.

Kaas P, Van Belle RA (1994) Monograph of Living Chitons: Mollusca: Polyplacophora). Volume 5. Suborder Ischnochitonina: Ischnochitonidae (concluded); Callistoplacinae, Mopalidae, additions to vols. 1–4. E.J. Brill, Leiden, the Netherlands, 402 pp.

Kaas P, Jones AM (1998) Class Polyplacophora: morphology and physiology. In: Beesley PL, Ross GJB, Wells A (Eds.) Mollusca: the southern synthesis part A. Fauna of Australia, CSIRO, Melbourne, Australia, 163–174.

Lopes BC, Silva EJ, Martins IX (2019) Looking for a home: the ecological association between the chitons Ischnochiton striolatus (Gray, 1828) and other molluscs species. Arquivo de Ciência do Mar (2): 52.

Matthews-Cascon H, Menezes TP (2019) Malacofauna associada a um recife de arenito no Nordeste do Brasil. Master's thesis, Universidade Federal do Ceará, Fortaleza, Brazil, 33 pp. http://repositorio.ufc.br/handle/riufc/42805. Accessed on: 2023-12-19.

Mendonça V, Vinagre C, Cabral H, Silva ACF (2015) Habitat use of inter-tidal chitons—role of colour polymorphism. Marine Ecology 36: 1098–1106. https://doi.org/10.1111/maec.12205

Menegotto A, Dambros CS, Netto SA (2019) The scale-dependent effect of environmental filters on species turnover and nestedness in an estuarine benthic community. Global Ecology and Biogeography 22: 639–647. https://doi.org/10.1002/ecy.2721

Morais JO, Freire GSS, Pinheiro LS, Souza MJN, Carvalho AM, Pessoa PRS, Oliveira SHM (2006) Ceará. In: Muehe D (Ed.) Caracterização fisiográfica e geoambiental da zona costeira do estado do ceará. Ministério do Ambiente, Rio de Janeiro, Brazil (1): 131–154.

Morais JO, Neto XAR, Pessoa PRS, Souza PL (2020) Morphological and sedimentary patterns of a semi-arid shelf, Northeast Brazil. Geo-Marine Letters 40: 835–842. https://doi.org/10.1007/s00367-019-00587-x

Nebra A, Alcaraz C, Caiola N, Muñoz-Camarillo G, Ibáñez C (2016) Benthic macrofaunal dynamics and environmental stress across a salt wedge Mediterranean estuary. Marine Environmental Research 117: 21–31. https://doi.org/10.1016/j.marenvres.2016.03.009

Righi G (1970) Mollusques Polyplacophores. Résultats scientifiques des campagnes de la "Calypso" (9). Campagne de *La Calypso* au large des côtes atlantiques de l'Amérique du Sud (1961–1962). Annales de l'Institut Océanographique de Monaco 47: 107–114.

Righi G (1971) Moluscos Poliplacóforos do Brasil. Papéis Avulsos de Zoologia 24: 123–146. https://doi.org/10.11606/0031-1049.1971.24.p123-146

Rocha-Barreira CA, Barros K, Matthews-Cascon H, Queiroz LR, da Silva AF, Barroso CX (2017) Ecology of mollusc communities in marine environments: central region of the semiarid coast of Brazil. In: Jenkins OP (Ed.). Advances in Animal Sciences and Zoology 10. Nova Science Publishers, New York, USA, 165–208.

Rostal DC, Simpson L (1988) The influence of salinity on the distribution of two Oregon chiton species (*Katharina tunicata* Wood and *Mopalia hindsii* Reeve). The Veliger 31: 120–126.

Schwabe E (2010) Illustrated summary of chiton terminology (Mollusca, Polyplacophora). Spixiana (2) 33: 171–194.

Shaw J, Maker G, Brooker L, Clode P (2009) The chiton fauna of the Swan River Estuary and their potential role as indicators of environmental contamination. The University of Western Australia, Australian (Data Manager) https://research-repository.uwa.edu.au/en/projects/the-chiton-fauna-of-the-swan-river-estuary-and-their-potential-ro-2. Accessed on: 2020-04-27.

Silva M (2000) Estuários—critérios para uma classificação ambiental. Revista Brasileira de Recursos Hídricos 5: 23–35. https://doi.org/10.21168/rbrh.v5n1.p23-35

Simone LRL, Jardim JA (2009) Classe Polyplacophora. In: Rios EC (Ed.) Seashells of Brazil. Second edition. Evangraf, Rio Grande, Brazil, 676 pp, 3–20.

Veiga MPT, Gutierre SMM, Castellano GC, Freire CA (2016) Tolerance of high and low salinity in the intertidal gastropod *Stramonita brasiliensis* (Muricidae): behaviour and maintenance of tissue water content. Journal of Molluscan Studies 82: 154–160. https://doi.org/10.1093/mollus/eyv044

Veras DRA, Martins IX, Matthews-Cascon H (2013) Mollusks: how are they arranged in the rocky intertidal zone? Iheringia, Série Zoologia 103: 97–103. https://doi.org/10.1590/S0073-47212013000200003

Vidy G (2000) Estuarine and mangrove systems and the nursery concept: which is which? The case of the Sine Saloum system (Senegal). Wetlands Ecology and Management 8: 37–51. https://doi.org/10.1023/A:1008402905530

Wolff WJ (2005) Non-indigenous marine and estuarine species in the Netherlands. Zoologische Mededelingen 79: 1–116.